Reprint from The Review of DIABETIC STUDIES

The Review Of
DIABETIC
STUDIES OPEN ACCESS

# Enhancing Healthcare Delivery and Institutional Performance, Through Integrated Nursing Leadership, Healthcare Administration, Health Information, Clinical Governance, and Quality Improvement Strategies.

**Dr Raid Mohammed Alawage[1], Abdulrhman Abdulrheem Khan[2], Mohammed Abed Albishri[3], Sanad Fahad Algethami[4], Mustafa Asaad Abu Hamamah[5], Tariq Sultan Alsafari[6], Nahed Meteb Althobaiti[7], Rasha Tariq Abdulljwad[8], Suleiman Mohammed Alawage[9], Faisal Shukri Refaei[10], Faisal Fahad Algethami[11], Mohammed Mansour Hameed Alqarafi[12], Farah Abdualrahman Alshubrmi[13], Abdulrahim Saleh Alahmadi[14], Khalid Ali A Alsharif[15]**

*[1]Health Administration, King Abdullah Medical City, Makkah, Saudi Arabia*
*[2]Health Administration Specialist, Al-Noor Specialist Hospital, Makkah, Saudi Arabia*
*[3]Hospital Management and Health Services, King Abdullah Medical City, Makkah, Saudi Arabia*
*4Specialist in Healthcare and Hospital Administration, King Abdullah Medical City, Makkah, Saudi Arabia*
*[5]Healthcare and Hospital Management, King Abdullah Medical City, Makkah, Saudi Arabia*
*[6]Health Administration Specialist, Asfan Health Center, Saudi Arabia*
*[7]Health Administration Specialist, King Abdullah Medical City, Makkah, Saudi Arabia*
*8Health Information Technician, King Fahad General Hospital, Saudi Arabia*
*[9]Nursing Technician, King Abdulaziz Hospital, Saudi Arabia*
*[10]Nursing Specialist, King Abdullah Medical City, Makkah, Saudi Arabia*
*[11]Nursing Specialist, King Abdulaziz Hospital, Makkah, Saudi Arabia*
*[12]Nursing Technician, Madinah Health Cluster, Madinah, Saudi Arabia*
*[13]Nursing Technician, King Salman Specialized Hospital, Saudi Arabia*
*[14]Nursing Technician, Psychiatric Hospital, King Salman bin Abdulaziz Medical City, Saudi Arabia*
*[15]Nurse, King Abdulaziz Hospital, Saudi Arabia*

**Abstract**

Background: A change in a hospital performance indicator may reflect different care, a different mix of patients, or a change in reporting. Objective: To examine how nursing leadership, administration, and health information can work through clinical governance to improve hospital care and discharge transitions. Methods: Targeted searches for this narrative review prioritized systematic reviews and comparative evaluations. Studies from Saudi Arabia informed the discussion of local implementation. Results: Higher registered nurse staffing was associated with better patient outcomes in longitudinal and comparative studies. Trials support tailored fall prevention and structured discharge support in selected settings. Better documentation and increased reporting can change measured performance without demonstrating a change in harm.Conclusion: The proposed model links frontline observations and verified data to a named decision maker and a scheduled review of outcomes and unintended effects. Its combined effect should be evaluated locally.

**Keywords:** nursing leadership; health administration; health information management; clinical governance; quality improvement; patient safety

## 1. Introduction

When reported inpatient falls increase, the committee reviewing them must decide whether to change care, change how it measures falls, or do both. Ward nurses can describe when and how patients mobilize. Administration can identify changes in staffing or bed occupancy. Health information staff can check the event definition and denominator. Clinical governance brings those observations together and gives someone the authority and resources to act. This is one way to apply Donabedian's distinction between the

conditions in which care is delivered, what staff do, and the outcomes patients experience. [1, 2]

Improvement work changes care and studies whether the change helps. These are related tasks: a ward may discover that a new handover prompt is usable in one week, while a hospital needs longer observation and a comparison to determine whether the prompt altered patient outcomes. This review examines how nursing leadership, healthcare administration, health information, and clinical governance can support each task in acute hospitals and during the transition out of hospital. Its focus is on patient safety, access and flow, workforce conditions, and measures that help leaders decide how to use limited resources. [3]

## 2. Methods

This narrative review drew on targeted searches for PubMed-indexed studies and journal reports available through November 16, 2026. Search concepts included hospital boards, safety culture, nursing leadership, nurse staffing and skill mix, boarding and discharge, data quality and coding, patient safety reporting, audit and feedback, and quality improvement evaluation. Additional searches sought hospital studies from Saudi Arabia to assess local relevance. Foundational publications were retained when they clarified constructs or methods.

Selection emphasized systematic reviews, randomized evaluations, and controlled or longitudinal studies for questions about effects. Descriptive studies informed understanding of implementation barriers or context. Findings were compared by design, setting, outcome definition, and risk of alternative explanations. No pooled estimates were calculated. The figures and indicator set are proposed applications of the synthesis; they are not results of a tested integrated intervention.

## 3. Results

### 3.1 Clinical Governance and Accountability

Clinical governance gives hospital leadership responsibility for the quality of care and a means to act when risks exceed the authority of an individual unit. A study of hospital boards in the United States and England linked attention to clinical quality and use of quality metrics with stronger management practices. Safety culture reviews report associations with patient outcomes, but differences in measurement and study design constrain causal interpretation. Accreditation may strengthen processes and compliance; evidence for consistent effects on mortality or healthcare-associated infections is less certain. A quality committee therefore needs to see whether accredited processes are followed at the bedside and whether patients benefit. [4-6]

Daily huddles can identify immediate hazards and route them to staff who can resolve them. Questions about recurrent risks, staffing capacity, or technology require a separate review with authority to allocate resources. For example, a ward manager can reorganize a fall risk handover; inadequate availability of mobility aids across wards may require a procurement decision. A governance decision should name the person responsible, specify the resources and escalation route, and set a date for reviewing what changed. [7]

Figure 1 shows how nursing, administration, and health information bring different evidence to one governance decision and how the result is reported back to the ward.





**Figure 1.** Proposed governance model for translating a care concern into action. The clinical governance forum assigns responsibility and resources; the improvement team returns results to frontline staff and decision makers.

### 3.2 Nursing Leadership, Staffing, and Patient Flow

Reviews link nursing leadership and favorable work environments with workforce and organizational outcomes. Patient outcomes have been studied less consistently, so the association is clearer for some staff outcomes than for specific changes in harm. Longitudinal studies of registered nurse staffing add evidence about patients' exposure to understaffed shifts. Recent cohorts also examine team composition and when understaffing occurs. Such findings support reviewing staffing by ward and shift, with attention to case mix and skill mix. A ward median used as a study comparator is not a recommended safe minimum. A prospective comparison of hospitals in Queensland provides policy evidence and illustrates the value of tracking outcomes as well as staffing. [8-13]

Staffing reviews should use hours actually worked and account for redeployments that cover a shortfall in one ward by reducing capacity in another. Administrators can pair this information with admissions, transfers, discharge readiness, and capacity at receiving units. In a French cohort of older adults, remaining in the emergency department overnight while awaiting admission was associated with higher in-hospital mortality. That finding identifies a vulnerable group and an important flow problem; it does not establish that every intervention to shorten boarding reduces deaths. Shorter waits are useful only when the receiving ward can deliver appropriate care. [14]

At discharge, nurses can verify that patients understand medication changes and warning signs, while administration can address access to follow-up and health information can link records across encounters. A randomized trial found lower 30-day hospital use with a discharge package that included a nurse discharge advocate and pharmacist follow-up. A Cochrane review concluded that individualized discharge planning probably produces small reductions in initial length of stay and readmission among older medical patients. For a hospital assessing its own program, any reduction in recorded readmissions should be interpreted in light of capture of visits to other institutions and changes in follow-up access. [15, 16]

### 3.3 Health Information and Performance Measurement

Every measure needs a defined numerator, eligible denominator, time window, data source, and owner. For example, falls per 1,000 patient-days allow comparison across months with different occupancy, provided fall definitions and patient-day rules stay stable. Event occurrence time should be distinguished from the time a report is entered. Missing records should be counted as missing, because their absence can create a false appearance of improvement. Data quality checks should compare source records with dashboard figures, test plausibility, and document changes in extraction or classification. [17, 18]

Coding and risk adjustment can change hospital comparisons. A decrease in mortality for patients coded with a principal diagnosis of pneumonia may reflect the reclassification of some severe cases as sepsis or respiratory failure. Use present-on-admission and relevant baseline information where available, and avoid adjusting away complications that care may have caused. Present comparable populations and stratify by ward or relevant patient group when a pooled result could obscure a problem. An increase in voluntary incident reports may reflect better reporting. Independent chart review or trigger-based detection helps distinguish reporting behavior from changes in harm. [19-22]

Barcode medication a dministration illustrates the link between electronic information and clinical work. One comparative study found fewer administration errors unrelated to timing after barcode-assisted electronic medication records were introduced. Observations in other hospitals documented workarounds when the system did not fit bedside practice, including scanning barcodes away from the patient. Leaders should pair electronic completion rates with observation of actual practice and give staff a route to report failures that require equipment or workflow changes. [23, 24] Table 1 lists measures for reviewing hospital performance across teams.

**Table 1.** Example measures for a joint hospital performance review

| Domain | Example measure | Interpretation check |
|---|---|---|
| Staffing | Worked registered nurse hours per patient-day | Examine ward and shift variation with patient acuity and redeployment. |
| Handover | Transfers with all required items documented ÷ eligible transfers | Sample records to confirm that receiving teams obtained the information. |
| Inpatient harm | Injurious falls per 1,000 patient-days | Reconcile reports with clinical notes; check for changes in injury definitions. |
| Patient flow | Median time from admission decision to arrival on the ward | Define eligible encounters and assess nights and peak periods separately. |
| Care transitions | Unplanned readmissions within 30 days ÷ eligible discharges | Specify planned exclusions and whether other hospitals are captured. |
| Balancing measures | Overtime hours per 100 hours worked; assisted mobilizations per eligible patient-day | Check for added staff burden and reduced patient mobility. |

### 3.4 Quality Improvement and Evaluation

The 2025 Cochrane review of audit and feedback found that it generally improved professional practice by a small amount, with substantial variation. Most evidence came from work on physicians' prescribing or test ordering. Hospital teams can use the findings to choose an actionable measure, discuss performance with recipients who can change it, and agree when to review progress. They should test whether improvement in the audited activity leads to the patient outcome they seek. [25]

Comparisons between settings show why the same label for an intervention can conceal different implementation and context. A Michigan collaborative reported a reduction in catheter-related bloodstream infections after a multifaceted intensive care intervention. In England, infection rates declined before a related national program and the decline did not accelerate after implementation. An international before-and-after surgical checklist study found reductions in complications and death; a population study after checklist adoption in Ontario found no significant reduction in those outcomes. These studies support checking baseline trends and how the work is carried out before attributing a change to an implementation program. [26-29]

A randomized evaluation of a tailored bedside fall prevention toolkit found fewer falls, but no significant effect on fall injuries. A later nonrandomized evaluation also found an association with fewer injurious falls. A ward testing such a program should track assisted mobility and access to toileting as well as falls, so it can see whether a lower fall rate was accompanied by a restriction on patient activity. [30, 31]

Figure 2 shows how a rise in reported falls can prompt a check of measurement and a review of care before a ward selects a change to test.





**Figure 2.** Response to an increase in reported falls. Measurement and care problems can coexist. Address immediate safety concerns while verifying the signal.

Plan–do–study–act cycles test changes on a small scale. Record the prediction, the group and period tested, any adaptations, and the next decision. Statistical process control charts flag unusual variation for investigation. An interrupted time series can estimate changes in level and trend while accounting for seasonality and correlation over time. A credible comparison series strengthens attribution. Reports should describe context, implementation, and evaluation. [32-35]Table 2 summarizes selected results and the limits of their interpretation.

**Table 2.** Selected hospital interventions and interpretation of their results

| Intervention and design | Reported result | Interpretation |
|---|---|---|
| Nurse staffing policy; prospective hospital comparison (Queensland). [13] | One fewer patient per nurse was associated with lower odds of 30-day mortality (adjusted OR 0.93; 95% CI, 0.86–0.99). | Policy exposure was not randomized. |
| Nurse and pharmacist discharge program; single-center randomized trial. [15] | Combined emergency visits and hospitalizations over 30 days: 0.314 vs 0.451 visits per person (IRR 0.695; 95% CI, 0.515–0.937). | Evaluated a package; individual effects were not isolated. |
| Barcode medication administration; before-and-after comparison. [23] | Administration errors unrelated to timing: 11.5% without vs 6.8% with barcode use. | Measured observed errors; mortality was not established. |
| Intensive care catheter safety program; collaborative cohort. [26] | Median infection rate decreased from 2.7 to 0 per 1,000 catheter-days at 3 months. | A median of zero does not mean zero infections across units. |
| Tailored bedside fall prevention; cluster randomized trial. [30] | Site-adjusted falls: 3.15 vs 4.18 per 1,000 patient-days with the toolkit vs usual care. | No significant reduction in fall injuries in this trial. |

CI, confidence interval; IRR, incidence rate ratio; OR, odds ratio.

## 4. Discussion

The evidence is most useful when hospitals match the strength of the claim to the design. A trial can establish an effect for the intervention and population studied. A comparison across hospitals can support an estimate but is still sensitive to differences in baseline staffing, patient mix, and concurrent programs. Local improvement reports can show that a process became feasible; a short before-and-after analysis provides weaker evidence that the process caused a change in harm. Governance bodies should use operational signals to investigate and act where needed while reserving claims of sustained benefit for stronger evaluations. [13, 27, 34]

Saudi surveys identify staffing and the response to error as issues for institutional review. A recent health cluster survey also identified hospital management support as an area for improvement. A national study relating safety cultures to reported sentinel events provides context for these findings, but the events reported and the events experienced are different measures. In Makkah, a before-and-after evaluation of a nurse-led catheter infection prevention initiative reported improvements. A Dhahran project reported better discharge experience after a local redesign. These evaluations identify practical ideas for testing elsewhere; they cannot establish nationally generalizable effects on harm, hospital use, or cost. [36-40]

A hospital or health cluster could begin with one priority and agree which decisions belong to the ward and which require executive support. For discharge, the ward may own communication with patients, administration may own appointment capacity, and the information team may check whether post-discharge visits are captured. A shared review can then compare patient outcomes, process completion, overtime, and access. Stratifying by service, language needs, or discharge destination can reveal benefits or delays hidden by an overall average. For staffing, the Queensland study estimated that avoided costs from shorter stays and fewer readmissions exceeded the added staffing cost. Local budget decisions still need to account for staffing prices, available capacity, and whether released bed-days can be used for other patients. [13]

This narrative review did not formally grade certainty or pool results. Selection may have missed relevant

interventions, and several domains rely on studies from particular systems and on outcomes measured in different ways. For example, staffing availability and cross-hospital access to follow-up data influence whether findings transfer. The proposed model is a way to assign responsibility and learn from local implementation; the effect of integrating all its components has not been established. Multisite evaluations with a prespecified outcome, a suitable comparison, and measures of costs and staff burden would help distinguish benefits of integration from concurrent changes.

## 5. Conclusion

Hospital performance review is most useful when it connects a patient care concern to reliable data, a decision maker with authority, and follow-up. Nursing leaders can identify problems in care, administration can address staffing and capacity, and health information teams can check how outcomes are measured. Clinical governance can then assign action and require an evaluation that considers harms, access, and workload. Evidence supports several specific interventions; hospitals should test how the components perform together in their own settings.

## References

1. Donabedian A. The quality of care. How can it be assessed? JAMA [Internet]. 1988 [cited 2026 Nov 16];260(12):1743-8. Available from: https://doi.org/10.1001/jama.260.12.1743
2. Scally G, Donaldson LJ. The NHS's 50 anniversary. Clinical governance and the drive for quality improvement in the new NHS in England. BMJ [Internet]. 1998 [cited 2026 Nov 16];317(7150):61-5. Available from: https://doi.org/10.1136/bmj.317.7150.61
3. Batalden PB, Davidoff F. What is "quality improvement" and how can it transform healthcare? Qual Saf Health Care [Internet]. 2007 [cited 2026 Nov 16];16(1):2-3. Available from: https://doi.org/10.1136/qshc.2006.022046
4. Tsai TC, Jha AK, Gawande AA, Huckman RS, Bloom N, Sadun R. Hospital board and management practices are strongly related to hospital performance on clinical quality metrics. Health Aff (Millwood) [Internet]. 2015 [cited 2026 Nov 16];34(8):1304-11. Available from: https://doi.org/10.1377/hlthaff.2014.1282
5. Braithwaite J, Herkes J, Ludlow K, Testa L, Lamprell G. Association between organisational and workplace cultures, and patient outcomes: systematic review. BMJ Open [Internet]. 2017 [cited 2026 Nov 16];7(11):e017708. Available from: https://doi.org/10.1136/bmjopen-2017-017708
6. Hussein M, Pavlova M, Ghalwash M, Groot W. The impact of hospital accreditation on the quality of healthcare: a systematic literature review. BMC Health Serv Res [Internet]. 2021 [cited 2026 Nov 16];21(1):1057. Available from: https://doi.org/10.1186/s12913-021-07097-6
7. Goldenhar LM, Brady PW, Sutcliffe KM, Muething SE. Huddling for high reliability and situation awareness. BMJ Qual Saf [Internet]. 2013 [cited 2026 Nov 16];22(11):899-906. Available from: https://doi.org/10.1136/bmjqs-2012-001467
8. Hult M, Terkamo-Moisio A, Kaakinen P, Karki S, Nurmeksela A, Palonen M, et al. Relationships between nursing leadership and organizational, staff and patient outcomes: A systematic review of reviews. Nurs Open [Internet]. 2023 [cited 2026 Nov 16];10(9):5920-5936. Available from: https://doi.org/10.1002/nop2.1876
9. Lake ET, Sanders J, Duan R, Riman KA, Schoenauer KM, Chen Y. A Meta-Analysis of the Associations Between the Nurse Work Environment in Hospitals and 4 Sets of Outcomes. Med Care [Internet]. 2019 [cited 2026 Nov 16];57(5):353-361. Available from: https://doi.org/10.1097/MLR.0000000000001109
10. Needleman J, Buerhaus P, Pankratz VS, Leibson CL, Stevens SR, Harris M. Nurse staffing and inpatient hospital mortality. N Engl J Med [Internet]. 2011 [cited 2026 Nov 16];364(11):1037-45. Available from: https://doi.org/10.1056/NEJMsa1001025
11. Griffiths P, Saville C, Ball J, Culliford D, Jones J, Lambert F, et al. Nursing Team Composition and Mortality Following Acute Hospital Admission. JAMA Netw Open [Internet]. 2024 [cited 2026 Nov 16];7(8):e2428769. Available from: https://doi.org/10.1001/jamanetworkopen.2024.28769
12. Morioka N, Moriwaki M, Miyawaki A, Saville C, Fushimi K, Griffiths P. Hospital Nurse Understaffing and Patient Mortality, Readmission, and Length of Stay. JAMA Netw Open [Internet]. 2026 [cited 2026 Nov 16];9(2):e2558235. Available from: https://doi.org/10.1001/jamanetworkopen.2025.58235

13. McHugh MD, Aiken LH, Sloane DM, Windsor C, Douglas C, Yates P. Effects of nurse-to-patient ratio legislation on nurse staffing and patient mortality, readmissions, and length of stay: a prospective study in a panel of hospitals. Lancet [Internet]. 2021 [cited 2026 Nov 16];397(10288):1905-1913. Available from: https://doi.org/10.1016/S0140-6736(21)00768-6
14. Roussel M, Teissandier D, Yordanov Y, Balen F, Noizet M, Tazarourte K, et al. Overnight Stay in the Emergency Department and Mortality in Older Patients. JAMA Intern Med [Internet]. 2023 [cited 2026 Nov 16];183(12):1378-1385. Available from: https://doi.org/10.1001/jamainternmed.2023.5961
15. Jack BW, Chetty VK, Anthony D, Greenwald JL, Sanchez GM, Johnson AE, et al. A reengineered hospital discharge program to decrease rehospitalization: a randomized trial. Ann Intern Med [Internet]. 2009 [cited 2026 Nov 16];150(3):178-87. Available from: https://doi.org/10.7326/0003-4819-150-3-200902030-00007
16. Gonçalves-Bradley DC, Lannin NA, Clemson L, Cameron ID, Shepperd S. Discharge planning from hospital. Cochrane Database Syst Rev [Internet]. 2022 [cited 2026 Nov 16];(2):CD000313. Available from: https://doi.org/10.1002/14651858.CD000313.pub6
17. Weiskopf NG, Weng C. Methods and dimensions of electronic health record data quality assessment: enabling reuse for clinical research. J Am Med Inform Assoc [Internet]. 2013 [cited 2026 Nov 16];20(1):144-51. Available from: https://doi.org/10.1136/amiajnl-2011-000681
18. Kahn MG, Callahan TJ, Barnard J, Bauck AE, Brown J, Davidson BN, et al. A Harmonized Data Quality Assessment Terminology and Framework for the Secondary Use of Electronic Health Record Data. EGEMS (Wash DC) [Internet]. 2016 [cited 2026 Nov 16];4(1):1244. Available from: https://doi.org/10.13063/2327-9214.1244
19. Lindenauer PK, Lagu T, Shieh MS, Pekow PS, Rothberg MB. Association of diagnostic coding with trends in hospitalizations and mortality of patients with pneumonia, 2003-2009. JAMA [Internet]. 2012 [cited 2026 Nov 16];307(13):1405-13. Available from: https://doi.org/10.1001/jama.2012.384
20. Pine M, Jordan HS, Elixhauser A, Fry DE, Hoaglin DC, Jones B, et al. Enhancement of claims data to improve risk adjustment of hospital mortality. JAMA [Internet]. 2007 [cited 2026 Nov 16];297(1):71-6. Available from: https://doi.org/10.1001/jama.297.1.71
21. Sari AB, Sheldon TA, Cracknell A, Turnbull A. Sensitivity of routine system for reporting patient safety incidents in an NHS hospital: retrospective patient case note review. BMJ [Internet]. 2007 [cited 2026 Nov 16];334(7584):79. Available from: https://doi.org/10.1136/bmj.39031.507153.AE
22. Classen DC, Resar R, Griffin F, Federico F, Frankel T, Kimmel N, et al. 'Global trigger tool' shows that adverse events in hospitals may be ten times greater than previously measured. Health Aff (Millwood) [Internet]. 2011 [cited 2026 Nov 16];30(4):581-9. Available from: https://doi.org/10.1377/hlthaff.2011.0190
23. Poon EG, Keohane CA, Yoon CS, Ditmore M, Bane A, Levtzion-Korach O, et al. Effect of bar-code technology on the safety of medication administration. N Engl J Med [Internet]. 2010 [cited 2026 Nov 16];362(18):1698-707. Available from: https://doi.org/10.1056/NEJMsa0907115
24. Koppel R, Wetterneck T, Telles JL, Karsh BT. Workarounds to barcode medication administration systems: their occurrences, causes, and threats to patient safety. J Am Med Inform Assoc [Internet]. 2008 [cited 2026 Nov 16];15(4):408-23. Available from: https://doi.org/10.1197/jamia.M2616
25. Ivers N, Yogasingam S, Lacroix M, Brown KA, Antony J, Soobiah C, et al. Audit and feedback: effects on professional practice. Cochrane Database Syst Rev [Internet]. 2026 [cited 2026 Nov 16];(6):CD000259. Available from: https://doi.org/10.1002/14651858.CD000259.pub5
26. Pronovost P, Needham D, Berenholtz S, Sinopoli D, Chu H, Cosgrove S, et al. An intervention to decrease catheter-related bloodstream infections in the ICU. N Engl J Med [Internet]. 2006 [cited 2026 Nov 16];355(26):2725-32. Available from: https://doi.org/10.1056/NEJMoa061115
27. Bion J, Richardson A, Hibbert P, Beer J, Abrusci T, McCutcheon M, et al. 'Matching Michigan': a 2-year stepped interventional programme to minimise central venous catheter-blood stream infections in intensive care units in England. BMJ Qual Saf [Internet]. 2013 [cited 2026 Nov 16];22(2):110-23. Available from: https://doi.org/10.1136/bmjqs-2012-001325

28. Haynes AB, Weiser TG, Berry WR, Lipsitz SR, Breizat AH, Dellinger EP, et al. A surgical safety checklist to reduce morbidity and mortality in a global population. N Engl J Med [Internet]. 2009 [cited 2026 Nov 16];360(5):491-9. Available from: https://doi.org/10.1056/NEJMsa0810119
29. Urbach DR, Govindarajan A, Saskin R, Wilton AS, Baxter NN. Introduction of surgical safety checklists in Ontario, Canada. N Engl J Med [Internet]. 2014 [cited 2026 Nov 16];370(11):1029-38. Available from: https://doi.org/10.1056/NEJMsa1308261
30. Dykes PC, Carroll DL, Hurley A, Lipsitz S, Benoit A, Chang F, et al. Fall prevention in acute care hospitals: a randomized trial. JAMA [Internet]. 2010 [cited 2026 Nov 16];304(17):1912-8. Available from: https://doi.org/10.1001/jama.2010.1567
31. Dykes PC, Burns Z, Adelman J, Benneyan J, Bogaisky M, Carter E, et al. Evaluation of a Patient-Centered Fall-Prevention Tool Kit to Reduce Falls and Injuries: A Nonrandomized Controlled Trial. JAMA Netw Open [Internet]. 2020 [cited 2026 Nov 16];3(11):e2025889. Available from: https://doi.org/10.1001/jamanetworkopen.2020.25889
32. Taylor MJ, McNicholas C, Nicolay C, Darzi A, Bell D, Reed JE. Systematic review of the application of the plan-do-study-act method to improve quality in healthcare. BMJ Qual Saf [Internet]. 2014 [cited 2026 Nov 16];23(4):290-8. Available from: https://doi.org/10.1136/bmjqs-2013-001862
33. Benneyan JC, Lloyd RC, Plsek PE. Statistical process control as a tool for research and healthcare improvement. Qual Saf Health Care [Internet]. 2003 [cited 2026 Nov 16];12(6):458-64. Available from: https://doi.org/10.1136/qhc.12.6.458
34. Bernal JL, Cummins S, Gasparrini A. Interrupted time series regression for the evaluation of public health interventions: a tutorial. Int J Epidemiol [Internet]. 2017 [cited 2026 Nov 16];46(1):348-355. Available from: https://doi.org/10.1093/ije/dyw098
35. Ogrinc G, Davies L, Goodman D, Batalden P, Davidoff F, Stevens D. SQUIRE 2.0 (Standards for QUality Improvement Reporting Excellence): revised publication guidelines from a detailed consensus process. BMJ Qual Saf [Internet]. 2016 [cited 2026 Nov 16];25(12):986-992. Available from: https://doi.org/10.1136/bmjqs-2015-004411
36. Alaska YA, Alkutbe RB. What Do We Know About Patient Safety Culture in Saudi Arabia? A Descriptive Study. J Patient Saf [Internet]. 2023 [cited 2026 Nov 16];19(8):517-524. Available from: https://doi.org/10.1097/PTS.0000000000001165
37. Khalil AAA, Alqaser M, Barakat SW, Kaddoura MA, Ishaqui AA, Iqbal A, et al. Patient Safety Culture in a Saudi Health Cluster: A Cross-sectional Survey. J Patient Saf [Internet]. 2026 Oct 26 [cited 2026 Nov 16]. Online ahead of print. Available from: https://doi.org/10.1097/PTS.0000000000001603
38. Binkheder S, Alaska YA, Albaharnah A, AlSultan RK, Alqahtani NM, Amr AA, et al. The relationships between patient safety culture and sentinel events among hospitals in Saudi Arabia: a national descriptive study. BMC Health Serv Res [Internet]. 2023 [cited 2026 Nov 16];23(1):270. Available from: https://doi.org/10.1186/s12913-023-09205-0
39. Tayyib N, Alsolami F. Assessing the Impact of a Preventive Care Bundle and Nursing Empowerment on Catheter-Associated Urinary Tract Infection (CAUTI) Rates in Saudi Arabia: A Single-Arm Intervention Study. SAGE Open Nurs [Internet]. 2025 [cited 2026 Nov 16];11:23779608251399263. Available from: https://doi.org/10.1177/23779608251399263
40. Almohaisen N, Woodman A, Abid MH, AlGhazali OS, Alnowaiser N. Patient Experience at Discharge: A Quality Improvement Study in an Armed Forces Hospital, Saudi Arabia. J Patient Exp [Internet]. 2025 [cited 2026 Nov 16];12:23743735251325507. Available from: https://doi.org/10.1177/23743735251325507